

AD-A038 945

WISCONSIN UNIV MADISON MATHEMATICS RESEARCH CENTER

F/G 12/1

ANOMALOUS CONVERGENCE OF A CONTINUED FRACTION FOR RATIOS OF KUM--ETC(U)

JAN 77 W GAUTSCHI

DAA629-75-C-0024

UNCLASSIFIED

MRC-TSR-1711

NL

3 OF 1  
AD  
A038945



END

DATE  
FILMED  
5-77

ADA 038945

MRC Technical Summary Report #1711

ANOMALOUS CONVERGENCE OF A  
CONTINUED FRACTION FOR RATIOS OF  
KUMMER FUNCTIONS

Walter Gautschi

12

Mathematics Research Center  
University of Wisconsin-Madison  
610 Walnut Street  
Madison, Wisconsin 53706

January 1977

Received December 28, 1976

See 1473

Approved for public release  
Distribution unlimited

DDC  
RECEIVED  
MAY 3 1977  
C

ADU NU.  
DDC FILE COPY

Sponsored by

U.S. Army Research Office  
P.O. Box 12211  
Research Triangle Park  
North Carolina 27709

and

National Science Foundation  
Washington, D. C. 20550

UNIVERSITY OF WISCONSIN - MADISON  
MATHEMATICS RESEARCH CENTER

ANOMALOUS CONVERGENCE OF A CONTINUED FRACTION  
FOR RATIOS OF KUMMER FUNCTIONS

Walter Gautschi

Technical Summary Report #1711  
January 1977

ABSTRACT

We exhibit a phenomenon of apparent convergence to the wrong limit in connection with a continued fraction of Perron for ratios of Kummer functions. The phenomenon is further illustrated in the special cases of Bessel functions and incomplete gamma functions.

AMS(MOS) Subject Classification 33A30, 33A40, 40A15, 65D20

Key Words: continued fractions, apparent convergence to the wrong limit, Bessel functions, incomplete gamma functions.

Work Unit Number 7 - Numerical Analysis.

Approved for	Write Name	
RTS	Dist. Status	
DAAG	UNCLASSIFIED	
	DISTRIBUTION	
BY	DISTRIBUTION/AVAILABILITY DATA	
Dist.	AVAIL. AND OR SPECIAL	
		A

# ANOMALOUS CONVERGENCE OF A CONTINUED FRACTION FOR RATIOS OF KUMMER FUNCTIONS

Walter Gautschi

1. Introduction. From the differential equation satisfied by Kummer's function  $M(a, b; z) = 1 + \frac{a}{b} \frac{z}{1!} + \frac{a(a+1)}{b(b+1)} \frac{z^2}{2!} + \dots$ , Perron [4, p. 278] develops the following continued fraction,

$$(1.1) \quad \frac{zM'(a, b; z)}{M(a, b; z)} = \frac{az}{b-z+} \frac{(a+1)z}{b+1-z+} \frac{(a+2)z}{b+2-z+} \dots, \quad b \neq 0, -1, -2, \dots,$$

where  $M'(a, b; z) = (d/dz)M(a, b; z) = (a/b)M(a+1, b+1; z)$ . While the continued fraction converges for any complex  $z$  not a zero of  $M(a, b; z)$ , the convergence behavior can be extremely deceptive, when  $|z| \gg \max(|a|, |b|)$ , particularly if  $\operatorname{Re} z > 0$ . The point is illustrated by concrete examples involving Bessel and incomplete gamma functions.

## 2. The phenomenon of apparent convergence to the wrong limit.

We assume, for simplicity, that  $a \neq 0$  and  $b-z \neq 0, -1, -2, \dots$ . Equation

(1.1) can then be written in the form

$$(2.1) \quad \frac{b-z}{a} \frac{M'(a, b; z)}{M(a, b; z)} = \frac{1}{1+} \frac{a_1}{1+} \frac{a_2}{1+} \dots,$$

where

$$(2.2) \quad a_k = \frac{(a+k)z}{(b-z+k-1)(b-z+k)}, \quad k = 1, 2, 3, \dots$$

Alternatively (cf., e.g., Wall [6, p. 17ff]),

$$(2.3) \quad \frac{b-z}{a} \frac{M'(a, b; z)}{M(a, b; z)} = \sum_{k=0}^{\infty} p_k,$$

where

$$p_0 = 1, \quad p_k = p_1 p_2 \dots p_k, \quad k = 1, 2, 3, \dots,$$

---

Sponsored by the United States Army under Contract No. DAAG29-75-C-0024 and The National Science Foundation under grant MCS 76-00842.

and

$$(2.4) \quad \rho_0 = 0, \quad \rho_k = \frac{-a_k(1+\rho_{k-1})}{1+a_k(1+\rho_{k-1})}, \quad k = 1, 2, 3, \dots$$

The infinite series in (2.3) represents the continued fraction in (2.1) in the sense that the  $n$ -th partial sum of the former is equal to the  $n$ -th convergent of the latter,  $n = 1, 2, 3, \dots$

Evidently, the terms  $p_k$  in the series of (2.3) decrease or increase in absolute value according to as  $|\rho_k| < 1$  or  $|\rho_k| > 1$ , respectively. It is useful, therefore, to examine the behavior of  $|\rho_k|$  as a function of  $k$ .

Assuming  $|\rho_{k-1}| < 1$ , then  $|\rho_k| < 1$  certainly if  $|a_k| \leq \frac{1}{4}$ . On the other hand, by (2.2), if  $|z| > |b| + k$ , then  $|a_k| < (|a| + k)|z|(|z| - |b| - k)^{-2}$ , and an elementary calculation shows the upper bound for  $|a_k|$  to be  $\leq \frac{1}{4}$  if

$$(2.5) \quad |z| \geq 2(|b| + 2|a| + 3k).$$

It follows that (2.5), together with  $|\rho_{k-1}| < 1$ , implies  $|\rho_k| < 1$ . Since, initially,  $\rho_0 = 0$ , we obtain by induction that  $|\rho_k| < 1$  for all  $k$  satisfying (2.5).

If  $|z|$  is large, we see that the terms  $p_k$  in (2.3) must decrease initially, the rate of decrease being greater the larger  $|z|$ . The continued fraction in (2.1) then gives the appearance of converging rapidly to a value of the order of magnitude 1, yielding for  $M'(a, b; z)/M(a, b; z)$  a value approximately equal to  $-a/z$ . This is obviously the wrong answer, if  $\text{Re } z > 0$ . Indeed, from known asymptotic formulas [5, Eq. 13.1.4],

$$(2.6) \quad \frac{M'(a, b; z)}{M(a, b; z)} \sim 1 \quad \text{as } |z| \rightarrow \infty \quad \text{in } \text{Re } z > 0.$$

What is likely to happen, then, is that the terms  $p_k$ , after the initial descent, begin to increase again, and converge to zero only after reaching some peak values which are sufficiently large so as to contribute to a limit consistent with (2.6). It is only during the "final descent" of the terms  $p_k$  that the correct limit will be attained (assuming no rounding errors).

The phenomenon of apparent convergence, while prevalent for  $\operatorname{Re} z > 0$  and  $|z|$  large, need not occur if  $\operatorname{Re} z < 0$ , since in this case [5, Eq. 13.1.5]

$$(2.7) \quad \frac{M'(a,b;z)}{M(a,b;z)} \sim -\frac{a}{z} \quad \text{as } |z| \rightarrow \infty \text{ in } \operatorname{Re} z < 0.$$

Nevertheless, we will see in examples that the phenomenon persists if

$$\frac{\pi}{2} \leq |\arg z| < \pi, \text{ albeit in a weakened form.}$$

3. The case of real  $z$ . It is instructive to examine in more detail the case of real arguments  $z$  and real parameters  $a, b$  satisfying  $0 < a+1$  if  $z > 0$ , and  $0 < a+1 \leq b$  if  $z < 0$ .

It will be convenient to introduce the quantities

$$(3.1) \quad \sigma_k = 1 + \rho_k, \quad k = 0, 1, 2, \dots,$$

for which the recursion (2.4) gives

$$(3.2) \quad \sigma_0 = 1, \quad \sigma_k = \frac{1}{1 + a_k \sigma_{k-1}}, \quad k = 1, 2, 3, \dots$$

3.1. The case  $z = x > 0$ ,  $a+1 > 0$ . We consider two subcases, (i)  $b-x > 0$ , (ii)  $b-x < 0$ .

In case (i), it follows from (2.2) that  $a_k > 0$  for all  $k \geq 1$ , hence from (3.2) that  $0 < \sigma_k < 1$  for all  $k \geq 1$ , and therefore from (3.1) that  $-1 < \rho_k < 0$ . We see that the terms  $p_k$  in (2.3) alternate in sign and decrease monotonically in modulus. In fact, since  $a_k \rightarrow 0$ , hence  $a_k \sigma_{k-1} \rightarrow 0$

as  $k \rightarrow \infty$ , we have  $\sigma_k \rightarrow 1$ , and so  $\rho_k \rightarrow 0$ , meaning that the series in (2.3) converges faster than any geometric series. Indeed,

$$(3.3) \quad \rho_k \sim -\frac{x}{k} \quad \text{as } k \rightarrow \infty,$$

as is easily verified.

In case (ii), there exists a unique integer  $k_0 \geq 1$  such that  $x-b < k_0 < x-b+1$ . Therefore,  $a_{k_0} < 0$ , while  $a_k > 0$  for all  $k \neq k_0$ . It follows as before that  $-1 < \rho_k < 0$  for  $k < k_0$ . If  $\sigma_{k_0} > 0$  (even though  $a_{k_0} < 0$ ), then  $-1 < \rho_k < 0$  also for all  $k \geq k_0$ , and we have the same alternating and supergeometric convergence behavior as in case (i). If, however,  $\sigma_{k_0} < 0$  (which will be the case if  $x$  is large), then  $|\rho_{k_0}| > 1$ . Since  $a_k > 0$  for  $k > k_0$ , the inequality  $|\rho_k| > 1$  will persist as long as  $\sigma_k$  remains negative. Eventually, however,  $\sigma_k$  has to turn positive (the series in (2.3) being convergent), and from this point on, all subsequent  $\sigma$ 's remain positive, hence the corresponding  $\rho$ 's less than one in modulus. Therefore, if  $x$  is large, the sequence  $\{|\rho_k|\}$  initially decreases, then increases, and finally decreases to zero at a supergeometric rate given by (3.3). The "dip-and-peak" effect is more pronounced, the larger  $x$ , and is what gives rise to the phenomenon of apparent convergence.

3.2. The case  $z = -x < 0$ ,  $0 < a+1 \leq b$ . This time,  $a_k < 0$  for all  $k \geq 1$ . Noting that the function  $x(b+x+k-1)^{-1}(b+x+k)^{-1}$  on  $0 \leq x < \infty$  assumes a unique maximum at  $(b+k-1)^{\frac{1}{2}}(b+k)^{\frac{1}{2}}$ , we find that

$$|a_k| \leq \frac{a+k}{(\sqrt{b+k-1} + \sqrt{b+k})^2}, \quad k \geq 1,$$

and thus, in particular,

$$|a_k| \leq \frac{a+k}{2b+2k-1+2\sqrt{b+k-1}\sqrt{b+k}} < \frac{a+k}{2b+2k-1+2(b+k-1)} < \frac{1}{4} \frac{a+k}{b+k-1}.$$

Since  $a+1 \leq b$ , it follows that

$$a_k < 0, \quad |a_k| < \frac{1}{4} \text{ for all } k \geq 1.$$

From this, and (3.2), we deduce inductively

$$1 < \sigma_k < \frac{2(k+1)}{k+2} \text{ for all } k \geq 1,$$

hence, in particular,  $0 < \rho_k < 1$  for all  $k \geq 1$ . The series in (2.3) is now a series of positive monotonically decreasing terms, and convergence thus monotone and, as before, supergeometric.

#### 4. Examples.

4.1. Bessel functions. We specialize (1.1) to  $a = \nu + \frac{1}{2}$ ,  $b = 2\nu + 1$ , where  $\nu \geq \frac{1}{2}$ , and use  $M(\nu + \frac{1}{2}, 2\nu + 1; z) = \Gamma(1+\nu) \exp(\frac{1}{2}z) (\frac{1}{4}z)^{-\nu} I_\nu(\frac{1}{2}z)$ , together with the differential-difference relation  $I'_\nu(z) = I_{\nu-1}(z) - \frac{\nu}{z} I_\nu(z)$ , to obtain

$$(4.1) \quad \frac{1}{2}z \left\{ \frac{I_{\nu-1}(\frac{1}{2}z)}{I_\nu(\frac{1}{2}z)} - \frac{4\nu}{z} + 1 \right\} = \frac{(\nu + \frac{1}{2})z}{2\nu+1-z} \frac{(\nu + \frac{3}{2})z}{2\nu+2-z} \frac{(\nu + \frac{5}{2})z}{2\nu+3-z} \cdots = \frac{(\nu + \frac{1}{2})z}{2\nu+1-z} \sum_{k=0}^{\infty} p_k.$$

In Figure 4.1, the moduli of the terms,  $|p_k|$ , are plotted in function of  $k$ , for  $\nu=1$ , and  $z = re^{i\varphi}$ ,  $r=10, 20, 40$ ,  $\varphi=0, \pi/8, 2\pi/8, \dots, \pi$ . The behavior of  $|p_k|$ , when  $r$  is fixed and  $\varphi$  varies between 0 and  $\pi/2$ , is almost identical and is represented by one curve in Figure 4.1. The dependence on  $\varphi$  is shown only in the case  $r = 40$ , but is analogous for the other values of  $r$ .




Figure 4.1 Anomalous convergence of (4.1) for  $v = 1$  and  $z = re^{i\varphi}$

It is seen, typically, that the terms  $|p_k|$  decrease rapidly at the beginning, then bottom out and rise to a sharp peak, before finally converging to zero. The dip of the curve, and the upswing that follows, are quite substantial if  $0 \leq \varphi \leq \pi/2$  and  $r$  is large, for reasons explained in Section 3.1, case (ii). As  $\varphi$  increases from  $\pi/2$  to  $\pi$ , the peaking of the curves gradually weakens and finally disappears when  $\varphi = \pi$ . A similar attenuation takes place upon increasing the value of  $\nu$ , as is to be expected from the discussion in Section 3.1, case (i).

The seriousness of the convergence anomaly can be seen, e.g., in the case  $\nu = 1$ ,  $r = 40$ ,  $0 \leq \varphi \leq \pi/2$ . If we require ten decimal digit accuracy, we will attain it at about  $k = 15$  and retain it through about  $k = 60$ , the partial sums in (4.1) all having the same value to ten decimal digits in the range  $15 \leq k \leq 60$ . This "apparent limit", of course, is totally incorrect, as the main contribution to the series comes from the few terms around  $k = 100$ . The situation is aggravated by the fact that the numerical process of generating the terms  $p_k$  is accompanied by a substantial loss of accuracy during the upswing of the curve, amounting to a loss of about 16 digits when  $r = 40$ . A further complication is the apparent lack of warning signals: Known a-posteriori error estimates (see, e.g., [3]) either do not apply, or seem to apply only in the region of "final descent".

On the other hand, the convergence behavior of the continued fraction in (4.1) is quite acceptable when  $\varphi = \pi$ , i.e.,  $z = -x$ ,  $x > 0$ , in which case (4.1) can be given the form

$$(4.2) \quad \frac{1}{2} x \left[ \frac{I_{\nu-1}(\frac{1}{2} x)}{I_{\nu}(\frac{1}{2} x)} - \frac{4\nu}{x} - 1 \right] = \frac{-(\nu + \frac{1}{2})x}{2\nu+1+x-} \frac{(\nu + \frac{3}{2})x}{2\nu+2+x-} \frac{(\nu + \frac{5}{2})x}{2\nu+3+x-} \cdots$$

Convergence is more rapid the larger  $x$  and/or  $\nu$ . The use of this continued fraction, in combination with Gauss' continued fraction, is further discussed in [2].

4.2. Incomplete gamma function. We have  $M(a, a+1; z) = a(-z)^{-a} \gamma(a, -z)$ , where  $\gamma(a, \cdot)$  denotes the incomplete gamma function. Noting that  $M'(a, a+1; z) = \frac{a}{a+1} M(a+1, a+2; z)$ , Equation (1.1) now takes the form

$$(4.3) \quad \frac{\gamma(a+1, -z)}{\gamma(a, -z)} = \frac{-az}{a+1-z+} \frac{(a+1)z}{a+2-z+} \frac{(a+2)z}{a+3-z+} \cdots$$

The convergence behavior of (4.3) for  $a > 0$  appears to be quite analogous to that of (4.1), exhibiting the phenomenon of apparent convergence for  $z$  large in the complex plane cut along the negative real axis. Along the negative real axis, we have monotone convergence, if  $a > -1$ , according to the results of Section 3.2. In this case,

$$(4.4) \quad \frac{\gamma(a+1, x)}{\gamma(a, x)} = \frac{ax}{a+1+x-} \frac{(a+1)x}{a+2+x-} \frac{(a+2)x}{a+3+x-} \cdots, \quad x > 0.$$

We may combine this with  $\gamma(a+1, x) = a\gamma(a, x) - x^a e^{-x}$  to obtain

$$(4.5) \quad x^{-a} e^x \gamma(a, x) = \frac{1}{a-} \frac{ax}{a+1+x-} \frac{(a+1)x}{a+2+x-} \frac{(a+2)x}{a+3+x-} \cdots, \quad x > 0.$$

The use of this continued fraction, in combination with other methods, to evaluate incomplete gamma functions is discussed in [1].

Department of Computer Sciences  
Purdue University  
Lafayette, Indiana 47907

and Mathematics Research Center  
University of Wisconsin-Madison  
Madison, Wisconsin 53706

### References

1. W. Gautschi, An evaluation procedure for incomplete gamma functions, to be published.
2. W. Gautschi and J. Slavik, On the computation of modified Bessel functions, to be published.
3. W. B. Jones, Analysis of truncation error of approximations based on the Padé table and continued fractions, Rocky Mountain J. Math. 4 (1974), 241-250.
4. O. Perron, Die Lehre von den Kettenbrüchen, Vol. II, 3<sup>rd</sup> ed., B. G. Teubner, Stuttgart, 1957.
5. L. J. Slater, Confluent hypergeometric functions, in: Handbook of Mathematical Functions (M. Abramowitz & I. A. Stegun, eds.), pp. 503-535. Nat. Bur. Standards Appl. Math. Series, Vol. 55 (1964).
6. H. S. Wall, Analytic Theory of Continued Fractions, D. van Nostrand Co., New York, 1948. {Reprinted in 1967 by Chelsea Publ. Co., Bronx, N.Y. }

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 1711	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER (9) Technical
4. TITLE (and Subtitle) (6) ANOMALOUS CONVERGENCE OF A CONTINUED FRACTION FOR RATIOS OF KUMMER FUNCTIONS		5. TYPE OF REPORT & PERIOD COVERED Summary Report, no specific reporting period
7. AUTHOR(s) (10) Walter Gautschi		8. CONTRACT OR GRANT NUMBER(s) (15) DAAG29-75-C-0024, NSF-MCS-76-00842
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of Wisconsin 610 Walnut Street Madison, Wisconsin 53706		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 7 Numerical Analysis
11. CONTROLLING OFFICE NAME AND ADDRESS See Item 18		12. REPORT DATE (11) January 1977
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) (12) 12p.		13. NUMBER OF PAGES 9
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited. (14) MRC-TSR-1711		15. SECURITY CLASS. (of this report) UNCLASSIFIED
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES U.S. Army Research Office and National Science Foundation P.O. Box 12211 Washington, D. C. 20550 Research Triangle Park, North Carolina 27709		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Continued fractions, apparent convergence to the wrong limit, Bessel functions, incomplete gamma functions		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) We exhibit a phenomenon of apparent convergence to the wrong limit, in connection with a continued fraction of Perron for ratios of Kummer functions. The phenomenon is further illustrated in the special cases of Bessel functions and incomplete gamma functions.		

221 200

4B